

# Discrete Optimization 2024 (EPFL): Problem set of week 12

May 22, 2024

1. Verify that  $\{(x, y) \in \mathbb{R}^2 \mid 5x^2 - 2xy + 2y^2 \leq 1\}$  is an ellipse and find its area.

**Solution:** We can write  $5x^2 - 2xy + 2y^2 = (x + y)^2 + (2x - y)^2$ . Therefore,  $T(x, y) = (x + y, 2x - y)$  takes the ellipse to the unit disc (of area  $\pi$ ). We see from here that  $T^{-1}$  takes the unit disc to the ellipse. Therefore, we have an ellipse of area  $\pi |\det(T^{-1})| = \pi \frac{1}{3}$ .

2. Let  $B$  be the unit ball in  $\mathbb{R}^3$  and let  $H$  be the hyperplane  $H = \{x + 3y + 2z = 0\}$ .

Consider the linear transformation

$$T = \begin{pmatrix} 1 & 1 & 2 \\ -1 & 0 & 1 \\ 2 & 1 & 0 \end{pmatrix}$$

What is the (two dimensional) area of  $T(B \cap H)$ ?

**Solution:** We find two linearly independent vectors in  $H$ :  $u = (2, 0, -1)$  and  $v = (3, -3, 0)$ . We calculate the area of the parallelogram they generate (equal to  $\sqrt{|u|^2|v|^2 - \langle u, v \rangle^2}$ ) and we get  $\sqrt{54}$ . Then we consider  $T(u) = (-1, 3, 4)$  and  $T(v) = (0, -3, 3)$  and compute the area of the parallelogram they generate:  $\sqrt{459}$ . Therefore, the area of  $T(B \cap H) = \frac{\sqrt{459}}{\sqrt{54}}\pi = \frac{\sqrt{17}}{\sqrt{2}}\pi$ .

3. Let  $E \subset \mathbb{R}^3$  be the ellipsoid  $E = \{(x, y, z) \mid \frac{(x+2y-z)^2}{4} + \frac{(y-x-z)^2}{9} + \frac{(2x-y-2z)^2}{25} \leq 2\}$ . What is the volume of  $E$ ?

**Solution:** We observe that if we consider any point  $(x, y, z)$  in  $E$  and then apply the linear transformation  $T(x, y, z) = ((x + 2y - z)/2, (y -$

$x - z)/3, (2x - y - 2z)/5$ , then  $(x, y, z)$  goes to a point in the unit ball of radius  $\sqrt{2}$  whose volume is  $(4/3)\pi(\sqrt{2})^3$ . The matrix representing  $T$  is the matrix  $\begin{pmatrix} 1/2 & 2/2 & -1/2 \\ -1/3 & 1/3 & -1/3 \\ 2/5 & -1/5 & -2/5 \end{pmatrix}$ . We calculate its determinant and find it is equal to  $-1/3$ . It follows now that  $1/3$  times the volume of  $E$  is equal to  $(4/3)\pi(\sqrt{2})^3$ . Therefore, the volume of  $E$  is equal to  $8\sqrt{2}\pi$ .

4. Let  $E \subset \mathbb{R}^3$  be the ellipsoid  $E = \{(x, y, z) \mid x^2 + \frac{y^2}{4} + \frac{z^2}{9} \leq 1\}$ .

Let  $H^+$  be the half-space  $H^+ = \{(x, y, z) \mid x + y + z \geq 0\}$ .

Find an ellipsoid  $E'$  such that  $E' \supset E \cap H^+$  and  $Vol(E') \leq Vol(E)e^{-1/(2(3+1))}$ .

**Solution:** The linear transformation  $T(x, y, z) = (x, \frac{y}{2}, \frac{z}{3})$  takes the ellipsoid  $E$  to the unit ball  $B$ . Notice that  $T$  takes  $H^+$  to the half-space  $G^+ = \{x + 2y + 3z \geq 0\}$ . Now we find a linear transformation  $R$  that will rotate the space about the origin such that  $B$  will remain fixed but  $G^+$  will go to the half space  $\{x \geq 0\}$ . This part is technical. One elegant way to do this is to take the vector  $u = \frac{1}{\sqrt{14}}(1, 2, 3)$  and complete it to an orthonormal basis: take also the vector  $v = \frac{1}{\sqrt{5}}(-2, 1, 0)$  and the vector  $w$  perpendicular to both  $u$  and  $v$  (you need to find it). Then consider the linear transformation  $R$  that takes  $u, v, w$  to  $(1, 0, 0), (0, 1, 0),$  and  $(0, 0, 1)$ . This is the inverse of the matrix whose columns are  $u, v, w$ .

The linear transformation  $R(T(\cdot))$  takes  $E \cap H^+$  to the half ball  $B \cap \{x \geq 0\}$ . By the half-ball theorem that we proved in class, the ellipsoid  $F = \{\frac{16}{9}(x - \frac{1}{4})^2 + \frac{8}{9}y^2 + \frac{8}{9}z^2 \leq 1\}$  contains the half ball  $B \cap \{x \geq 0\}$  and we also have  $Vol(F) \leq e^{\frac{-1}{2(3+1)}} Vol(B)$ .

If we now apply  $T^{-1}(R^{-1}(\cdot))$  on  $F$  we get the desired ellipsoid  $E'$ . One can get from here the algebraic description of  $E'$ . Technically, we need to replace  $x, y,$  and  $z$  in the algebraic description of  $F$  by  $R(T(x, y, z))$ .